Regional patterns of sea level change in the German North Sea in a worldwide context

UNIVERSITÄT SIEGEN

University of Siegen | Research Institute for Water and Environment | Dept. of Hydraulic Engineering Paul-Bonatz-Str. 9-11 | 57076 Siegen | Germany | phone +49 271 740 3462 | e-mail: thomas.wahl@uni-siegen.de

I. Introduction

Sea Level Rise (SLR) is one of the major consequences we are facing in times of a warming climate and it is obvious that a higher sea level influences the heights of occurring storm surges and thus results in a higher risk of inundation for the affected coastal areas.

Therefore, regional and global sea level rise are subjects to many recent scientific publications. In contrast, the mean sea 50% level (MSL) and its variability over the last centuries in the German North Sea area have not been analysed in detail up to now. A methodology to analyse observed sea level rise in the German Bight, the shallow south-eastern part of the North Sea, is presented. The contribution focuses on the description of the methods used to generate and analyse high quality mean sea level time series. A virtual station time series for the German Bight is estimated, analysed in detail by applying parametric and non-parametric fitting approaches and compared to two global sea level reconstructions (Church and White, 2006; Jevrejeva et al., 2008).



II. Data and Methods

The data sets of 13 tide gauges covering the entire German North Sea coastline are considered for the analyses (see Fig. 1). Figure 2 shows that data sets with a resolution in time of at least one hour are available since the end of the last century for most of the tide gauges. To combine the MSL values with the mean tide level (MTL) values, estimated from tidal high and low waters, the k-factor method is used (Wahl et al., 2010). Different tests are applied (Kolmogorov-Smirnov-Test, Mann-Kendall-Test, Sliding-Window-Test) to prove stationarity of the k-factor time series.

List		$k_{1937-1997} = 0,4539$
Hörnum		$k_{1937-1976} = 0,4442$ $k_{1977-1999} = 0,4575$
Wyk		$k_{1953-1994} = 0,4487$
Dagebüll		k ₁₉₃₆₋₁₉₉₉ = 0,4363
Wittdün		k ₁₉₃₆₋₁₉₉₉ = 0,4516
Husum		k ₁₉₃₆₋₁₉₉₉ = 0,4430
Helgoland	-	k ₁₉₉₀₋₁₉₉₆ = 0,4782
Cuxhaven	k ₁₈₄₃₋₁₉₁₇ = 0,4703	
Bremerhaven		k ₁₉₃₆₋₁₉₉₄ = 0,4493
Lt. Alte Weser		k ₁₉₀₁₋₁₉₉₄ = 0,4800
Wilhelmshaven		$k_{1881-1950} = 0,4597$ $k_{1975-1999} = 0,4670$
Norderney		k ₁₉₀₁₋₁₉₉₉ = 0,4874
Emden		k ₁₉₅₀₋₁₉₉₉ = 0,4286
ŀ	= High and low waters available	= High resolution data available (at least hourly data)
184	3 1875 1900	1925 1950 1975 2000

Figure 2. Available data sets and the estimated k-factors for different time periods.

Afterwards, a virtual station time series for the German Bight is estimated by integrating the averaged rates of SLR per year of the different gauges. Parametric fitting approaches as well as non-parametric data adaptive filters, such as Singular System Analysis (SSA), are applied to the resulting time series. For padding non-stationary sea level time series, an advanced approach named Monte-Carlo autoregressive padding (MCAP) is used. This approach allows the specification of a kind of uncertainties of the behaviour of smoothed time series near the boundaries. Finally, the reconstructed time series for the German Bight is connected to global sea level reconstructions.

Acknowledgments:	This	his is a German				Coastal Engin			neering	Research	**	Bundesministeri für Bildung
	Cour	ICII	(NFN)	pro	Jeci,	IUſ	idea	by	Ine	German		und Forschung
Federal Ministry of Education and Research (BMBF).												

<u>Thomas Wahl, Torsten Frank and Jürgen Jensen</u>

Kuratorium für Forschung im Küsteningenieurwesen

III. Results

Figure 3 (top) shows the estimated virtual station time series of the German Bight. This relative mean sea level time series has a linear trend of 2.01 ± 0.08 mm/a for the entire period since 1843. The estimated trend is 1.68 ± 0.14 mm/a for the period since 1901, 1.96 ± 0.26 for the period since 1937, 2.14 ± 0.39 mm/a for the period since 1951 and 3.60 ± 0.74 mm/a for the reduced period since 1971 (quoted errors are 1- σ standard errors).

In addition, figure 3 (top) shows the results from analysing the virtual station time series using SSA with an embedding dimension of 15 years and 75,000 MCAP simulations to achieve stable results for the reconstruction near the boundaries. Figure 3 (bottom) shows the first deviation of the reconstruction providing the best fit for the observed data. The results indicate that an accerleration of SLR took place at the end of the 19th century and was followed by a deceleration. Another period of accelerated SLR obviously started around 1970 and has declined in recent years. Actual rates of relative SLR in the German Bight are in the order of 4 mm/a (note that the uncertainties increase near the boundaries).



Figure 3. Virtual station for the German Bight and results from SSA reconstruction using the MCAP approach.

Figure 4 shows the results from estimating running linear trends for different time spans of the virtual station time series. The results for 30-, 40- and 50-a time spans approve the existance of a period of acceleration at the end of the 19th century and another one with its starting point around 1970. A window length of 20-a seems to be to short to achieve meaningful results for the underlying time series, showing a high variance.



References: Church, J. A., and N. J. White (2006): A 20th century acceleration in global sea-level rise, Geophys. Res. Lett., 33, L01602, doi:10.1029/2005GL024826. Jevrejeva, S., J. C. Moore, A. Grinsted, and P. L. Woodworth (2008): Recent global sea level acceleration started over 200 years ago?, Geophys. Res. Lett., 35, L08715, doi:10.1029/2008GL033611. Wahl, T., Jensen, J., and Frank, T. (2010): On analysing sea level rise in the German Bight since 1844, Nat. Hazards Earth Syst. Sci., 10, 171-179, 2010.

Figure 5 shows the results from comparing the virtual station time series of the German Bight with two global sea level reconstructions (Church and White, 2006; Jevrejeva et al., 2008). The reconstruction by Jevrejeva et al. (2008) is based on tide gauge data and provides MSL values for the period of 1700 to 2002. The reconstruction by Chuch and White (2006) considers tide gauge data as well as altimetry data and provides MSL values for the period of 1870 to 2007. Both reconstructions were GIA corrected. They are also analysed using SSA with an emb. dim. of 15 years in combination with MCAP. Figure 5 (top) shows the differences between the rates of SLR estimated from the GIA corrected reconstruction for the German Bight and the global reconstructions. Higher rates in the German Bight are observed for a period around 1850, a period around 1900 and the period covering the last 10 to 15 years.

The results clearly reveal the existance of different patterns of sea level change, which is approved by estimating 20-a running correlation coefficents (Figure 5, bottom). Insignificant correlations are present for most of the periods of the considered time span (significance levels were estimated by applying t-test statistics).



Figure 5. Difference of rate of SLR and 20-a running correlation coefficients between the virtual station for the German Bight and two global sea level reconstructions.

IV. Conclusions and Outline

- one covering the last decades.
- observations.

- more detailed and on different time scales.

Research Institute for And Environment And Environment And Environment Dept. of Hydraulic Engineering

• The long-term trends estimated from the reconstructed relative mean sea level time series of the German Bight are similar to those found from other regional or global studies.

• An accelerated SLR in the German Bight is detected for a period around 1900 and for another

• The estimated high recent rates of relative SLR are not unusual considering the last 165 years of

• The comparison of the reconstruction of the German Bight with two global sea level reconstructions clearly reveals the existance of different patterns of sea level change.

• This confirms the necessity of regional climate and SLR scenarios for coastal planning purposes. • More powerful methods have to be applied to analyse the available sea level reconstructions